



STEPHAN BAKER
MD, MBA, FACS



PLASTIC SURGERY OF THE FACE, BREAST AND BODY
CERTIFIED BY THE AMERICAN BOARD OF PLASTIC SURGERY

THE OFFICES AT MERRICK PARK 4425 PONCE DE LEON BLVD. SUITE 200 CORAL GABLES, FLORIDA 33146 - 305.381.8837 - drbaker.com

The following medications should be discontinued two weeks before surgery to minimize the chances for unnecessary bleeding:

Actron	Clinoril	Monacet	Robaxisal
Actab	Comeback	Motrin	Rufen
Advil	Congespirin	Multivitamins	Red Wine
Aggrenox	Cope	Nalfon	Sine-Aid
Aleve	Coricidin	Naprosyn	Sine-Off
Alka-Seltzer	Cosprin	Naproxen	Sodium
Alteplase	Coumadin	Norgesic	Salicylate
Aluprin		Norgesin	Sominex
Anacin	Darvon	Nuprin	SK-65
Anaprox	Compound	Nyquil	Stanback
Ansaid	Daypro	Nytol	Stendin
APC	(Oxaprozan)		Stero-Darvon
Arthropan	Dong Quai	Orudis	with A.S.A.
ASA	Diflunazil	Oruvail	Supac
Ascodeen-30	Dipyramidole	Oxycodone	Sure Sleep
Ascriptin	Diclofenac	Oxyphenbutazone	Synalgos
Aspergum	Disalcid		
Aspirin	Doan's Pills	Pabalate	Talwin
Axotal	Dolar	Pabirin	Tandearil
	Dolobid	Pamprin	Ticlid
Bayer Aspirin	Dristan	Panalgesic	(Ticlopidine)
Bayer Time Release	Duradyne	Pepto-Bismol	Tolectin
BC Tablet and	Duragesic	Percodan	Tolmetin
Powder	Duramed	Persantine	Toradol
Buff-A-Comp		Persistin	Triaminicin
Buffadyne	Easprin	Phenaphen	Trilisate
Bufferin	Ecotrin	Phenylbutazone	
Butalbital	Empirin	PMS	Vanquish
Butazolidin	Emprazil	Ponstel	Vitamin C
	Encaprin	Premsyn	Vitamin E
Cataflam	Equagesic	Propoxythene	Voltaren
Cama-Inlay Tabs	Etodolac	Plavix	
CamaArthritis Pain	Excedrin		Warfarin
Reliever	Extra Strength	Quiet World	
Chamomile		Resolve	Zomax
Cheracol Capsules			Zorpri